

A search for astrophysical burst signals at the Sudbury Neutrino Observatory

Article (Accepted Version)

Peeters, S J M and The SNO Collaboration, (2015) A search for astrophysical burst signals at the Sudbury Neutrino Observatory. *Astroparticle Physics*, 55. pp. 1-7. ISSN 0927-6505

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/66810/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

A Search for Astrophysical Burst Signals at the Sudbury Neutrino Observatory

B. Aharmim⁶, S. N. Ahmed¹⁴, A. E. Anthony^{17,a}, N. Barros⁸, E. W. Beier¹³, A. Bellerive⁴, B. Beltran¹, M. Bergevin^{7,5,b}, S. D. Biller¹², K. Boudjemline^{4,14}, M. G. Boulay¹⁴, B. Cai¹⁴, Y. D. Chan⁷, D. Chauhan⁶, M. Chen¹⁴, B. T. Cleveland¹², G. A. Cox^{19,c}, X. Dai^{14,12,4}, H. Deng¹³, J. A. Detwiler⁷, M. DiMarco¹⁴, M. D. Diamond⁴, P. J. Doe¹⁹, G. Doucas¹², P.-L. Drouin⁴, F. A. Duncan^{16,14}, M. Dunford^{13,d}, E. D. Earle¹⁴, S. R. Elliott^{9,19}, H. C. Evans¹⁴, G. T. Ewan¹⁴, J. Farine^{6,4}, H. Fergani¹², F. Fleurot⁶, R. J. Ford^{16,14}, J. A. Formaggio^{11,19}, N. Gagnon^{19,9,7,12}, J. T. M. Goon¹⁰, K. Graham^{4,14}, E. Guillian¹⁴, S. Habib¹, R. L. Hahn³, A. L. Hallin¹, E. D. Hallman⁶, P. J. Harvey¹⁴, R. Hazama^{19,e}, W. J. Heintzelman¹³, J. Heise^{2,9,14,f}, R. L. Helmer¹⁸, A. Hime⁹, C. Howard¹, M. Huang^{17,6,g}, P. Jagam⁵, B. Jamieson^{2,h}, N. A. Jelley¹², M. Jerkins¹⁷, K. J. Keeter^{20,14}, J. R. Klein^{17,13}, L. L. Kormos¹⁴, M. Kos^{14,i}, C. Kraus^{14,6}, C. B. Krauss¹, A. Krueger⁶, T. Kutter¹⁰, C. C. M. Kyba^{13,j}, R. Lange³, J. Law⁵, I. T. Lawson^{16,5}, K. T. Lesko⁷, J. R. Leslie¹⁴, I. Levine⁴, J. C. Loach^{12,7,k}, R. MacLellan^{14,l}, S. Majerus¹², H. B. Mak¹⁴, J. Maneira⁸, R. Martin^{14,7}, N. McCauley^{13,12,m}, A. B. McDonald¹⁴, S. R. McGee¹⁹, M. L. Miller^{11,n}, B. Monreal^{11,o}, J. Monroe^{11,p}, B. G. Nickel⁵, A. J. Noble¹⁴, H. M. O’Keeffe¹², N. S. Oblath^{19,11}, R. W. Ollerhead⁴, G. D. Orebi Gann^{12,13,q}, S. M. Oser², R. A. Ott¹¹, S. J. M. Peeters^{12,r}, A. W. P. Poon⁷, G. Prior^{7,d}, S. D. Reitzner⁵, K. Rielage^{9,19}, B. C. Robertson¹⁴, R. G. H. Robertson¹⁹, M. H. Schwendener⁶, J. A. Secrest^{13,s}, S. R. Seibert^{17,9,13}, O. Simard⁴, J. J. Simpson⁵, D. Sinclair^{4,18}, P. Skensved¹⁴, T. J. Sonley^{11,t}, L. C. Stonehill^{9,19}, G. Tešić^{4,u}, N. Tolich¹⁹, T. Tsui², R. Van Berg¹³, B. A. VanDevender^{19,h}, C. J. Virtue⁶, B. L. Wall¹⁹, D. Waller⁴, H. Wan Chan Tseung^{12,19}, D. L. Wark^{15,v}, P. J. S. Watson⁴, J. Wendland², N. West¹², J. F. Wilkerson^{19,w}, J. R. Wilson^{12,x}, J. M. Wouters^{9,y}, A. Wright¹⁴, M. Yeh³, F. Zhang⁴, K. Zuber^{12,z}

SNO Collaboration

ABSTRACT

The Sudbury Neutrino Observatory (SNO) has confirmed the standard solar model and neutrino oscillations through the observation of neutrinos from the solar core. In this paper we present a search for neutrinos associated with sources other than the solar core, such as gamma-ray bursters and solar flares. We present a new method for looking for temporal coincidences between neutrino events and astrophysical bursts of widely varying intensity. No correlations were found between neutrinos detected in SNO and such astrophysical sources.

¹Department of Physics, University of Alberta, Edmonton, Alberta, T6G 2R3, Canada

²Department of Physics and Astronomy, University of British Columbia, Vancouver, BC V6T 1Z1, Canada

³Chemistry Department, Brookhaven National Labora-

tory, Upton, NY 11973-5000

⁴Ottawa-Carleton Institute for Physics, Department of Physics, Carleton University, Ottawa, Ontario K1S 5B6, Canada

⁵Physics Department, University of Guelph, Guelph,

Ontario N1G 2W1, Canada

⁶Department of Physics and Astronomy, Laurentian University, Sudbury, Ontario P3E 2C6, Canada

⁷Institute for Nuclear and Particle Astrophysics and Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720

⁸Laboratório de Instrumentação e Física Experimental de Partículas, Av. Elias Garcia 14, 1º, 1000-149 Lisboa, Portugal

⁹Los Alamos National Laboratory, Los Alamos, NM 87545

¹⁰Department of Physics and Astronomy, Louisiana State University, Baton Rouge, LA 70803

¹¹Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, MA 02139

¹²Department of Physics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

¹³Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104-6396

¹⁴Department of Physics, Queen's University, Kingston, Ontario K7L 3N6, Canada

¹⁵Rutherford Appleton Laboratory, Chilton, Didcot OX11 0QX, UK

¹⁶SNOLAB, Sudbury, ON P3Y 1M3, Canada

¹⁷Department of Physics, University of Texas at Austin, Austin, TX 78712-0264

¹⁸TRIUMF, 4004 Wesbrook Mall, Vancouver, BC V6T 2A3, Canada

¹⁹Center for Experimental Nuclear Physics and Astrophysics, and Department of Physics, University of Washington, Seattle, WA 98195

^aPresent address: Center for Astrophysics and Space Astronomy, University of Colorado, Boulder, CO 80309

^bPresent address: Department of Physics, University of California, Davis, CA

^cPresent address: Institut für Experimentelle Kernphysik, Karlsruher Institut für Technologie, Karlsruhe, Germany

^dPresent address: CERN, Geneva, Switzerland

^ePresent address: Department of Physics, Hiroshima University, Hiroshima, Japan

^fPresent address: Sanford Underground Research Laboratory, Lead, SD

^gPresent address: Center of Cosmology and Particle Astrophysics, National Taiwan University, Taiwan

^hPresent address: Department of Physics, University of Winnipeg, Winnipeg, MB R3B 2E9, Canada

ⁱPresent address: Pacific Northwest National Laboratory, Richland, WA

^jPresent address: Institute for Space Sciences, Freie Universität Berlin, Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Germany

^kPresent address: Department of Physics, Shanghai Jiaotong University, Shanghai, China

^lPresent address: Department of Physics and Astron-

1. Introduction

The Sudbury Neutrino Observatory (SNO) collaboration has looked for time-dependent anomalies due to both periodic variations (Aharmim et al. 2005, 2010) in neutrino flux and short bursts of neutrinos (Aharmim et al. 2011b). In the present study, we are specifically searching for neutrino events that are correlated to other known astrophysical events. There is a wide variety of potential astrophysical sources of neutrinos. In this paper we consider “burst” events, which include γ -ray bursters (GRB’s), solar flares, magnetars, and an intense burst observed in the Parkes radio telescope. These astrophysical burst events are short-lived and occur at random. It is expected that any related neutrino signal would be undetectable (Piran 2004; Bahcall 1988) and the present experimental constraints provide only limits (Hirata et al. 1988, 1990; Aglietta et al. 1991; Fukuda et al. 2002; Thrane et al. 2009).

Similar searches for neutrinos temporally cor-

omy, University of Alabama, Tuscaloosa, AL

^mPresent address: Department of Physics, University of Liverpool, Liverpool, UK

ⁿPresent address: Center for Experimental Nuclear Physics and Astrophysics, and Department of Physics, University of Washington, Seattle, WA 98195

^oPresent address: Dept. of Physics, University of California, Santa Barbara, CA

^pPresent address: Dept. of Physics, Royal Holloway University of London, Egham, Surrey, UK

^qPresent address: Physics Department, University of California at Berkeley, and Lawrence Berkeley National Laboratory, Berkeley, CA

^rPresent address: Department of Physics and Astronomy, University of Sussex, Brighton BN1 9QH, UK

^sPresent address: Dept. of Chemistry and Physics, Armstrong Atlantic State University, Savannah, GA

^tPresent address: Dept. of Physics, Queen’s University, Kingston, Ontario, Canada

^uPresent address: Physics Department, McGill University, Montreal, QC, Canada

^vAdditional Address: Imperial College, London SW7 2AZ, UK

^wPresent address: Department of Physics, University of North Carolina, Chapel Hill, NC

^xPresent address: Dept. of Physics, Queen Mary University, London, UK

^yDeceased

^zPresent address: Institut für Kern- und Teilchenphysik, Technische Universität Dresden, 01069 Dresden, Germany

related with astrophysical events have treated all bursts (regardless of their intensity) equally in terms of their potential for ν production (Fukuda et al. 2002), and/or examined only the highest-intensity burst(s) (Thrane et al. 2009). We introduce a novel technique, the Maximum Likelihood Burst Analysis (MLBA), which features some advantages over previous analysis techniques when searching data from neutrino detectors for temporal correlations with astrophysical burst events. The MLBA provides the versatility to deal with bursts whose intensities vary over many orders of magnitude, and to accommodate correlations between γ intensity and ν emission. This allows us to use the integrated intensity over a large number of bursts instead of only the most intense burst. The results from the MLBA can therefore be employed to test a variety of models in a straightforward manner and, as with previous analysis techniques, can be employed to set limits on the number of neutrino events correlated with the bursts – and hence indirectly set limits on the fluences.

We begin with a discussion of the SNO detector before discussing two types of astrophysical events, GRB's and solar flares, on which we have chosen to focus. In section 4, we present the MLBA method and motivation. Next, we examine solar flare and GRB observations in conjunction with the SNO neutrino data set, and give limits for the associated parameter for neutrino fluences from the burst events. We also show how this can be related to the neutrino fluence limits that have been typically presented in the literature. In the final section, we search for signals in the SNO data associated with two unusual isolated astrophysical events: the Parkes radio burst (Lorimer et al. 2007) and the SGR 1806-20 magnetar eruption (Palmer et al. 2005). In all cases, no significant correlations are seen between the SNO data and the astrophysical sources. However, limits on neutrino fluences from solar flares and GRB's are improved in the low-energy ν regime compared to previous analyses in the literature.

2. The SNO Detector

The SNO detector (Boger et al. 2000), shown schematically in Figure 1, consisted of an inner volume containing 10^6 kg of 99.92% isotopically pure heavy water ($^2\text{H}_2\text{O}$, hereafter referred




Fig. 1.— (Color online) Schematic diagram of the SNO detector. We used a coordinate system with the center of the detector as the origin, and z direction as vertically upward.

to as D_2O) within a 12 m diameter transparent acrylic vessel (AV). Over 7×10^6 kg of H_2O between the rock and the AV shielded the D_2O from external radioactive backgrounds. An array of 9456 inward-facing 20 cm Hamamatsu R1408 photomultiplier tubes (PMTs), installed on an 17.8 m diameter stainless steel geodesic structure (PSUP), detected Cherenkov radiation produced in both the D_2O and H_2O . The PMT thresholds were set to $1/4$ of the charge from a single photoelectron. The inner 1.7×10^6 kg of H_2O between the AV and the PSUP shielded the D_2O against radioactive backgrounds from the PSUP and PMTs.

The detector was located in Vale's Creighton mine ($46^\circ 28' 30''$ N latitude, $81^\circ 12' 04''$ W longitude) near Sudbury, Ontario, Canada, with the center of the detector at a depth of 2092 m (5890 ± 94 meters water equivalent). At this depth, the rate of cosmic-ray muons entering the detector was approximately three per hour. Ninety-one outward-facing PMTs attached to the PSUP detected cosmic-ray muons. An offline veto based on information from these PMTs significantly reduced cosmogenic backgrounds.

SNO detected low energy neutrinos through the following reactions:

- $\nu_x + d \rightarrow p + n + \nu_x$, Neutral current (NC),
- $\nu_e + d \rightarrow p + p + e^-$, Neutrino Charged current (CC),
- $\bar{\nu}_e + d \rightarrow n + n + e^+$, Anti-Neutrino Charged current (CC-anti),
- $\nu_x + e^- \rightarrow \nu_x + e^-$, Elastic scattering (ES),

where x implies electron, muon or tau neutrinos. The NC reaction is equally sensitive to all three active neutrino and anti-neutrino flavors. This reaction is sensitive down to neutrino energies of approximately 2.2 MeV, which gives us a lower threshold than many other experiments. The CC reaction is only sensitive to ν_e s, and the CC-anti reaction is only sensitive to $\bar{\nu}_e$ s. The ES reaction is sensitive to all neutrino and anti-neutrino flavors, but the cross-section for ν_e s ($\bar{\nu}_e$ s) is approximately six (three) times larger than that for the other flavors. For neutrino energies above approximately 6 MeV, the CC reaction has a cross-section more than ten times higher than the ES reaction, which provides increased sensitivity for an equivalent target exposure compared to experiments that are only sensitive to the ES reaction.

The recoil electrons from both the ES, CC, and CC-anti reactions were detected directly through their production of Cherenkov light. The total amount of light detected by the PMT array was correlated with the energy of the recoil electron.

The SNO detector operated in three distinct phases distinguished by how the neutrons from the NC interactions were detected. In the D₂O phase, the detected neutrons captured on deuterons in the D₂O releasing a single 6.25 MeV γ -ray, and it was the Cherenkov light of secondary Compton electrons or e^+e^- pairs that was detected. In the salt phase, 2×10^3 kg of NaCl were added to the D₂O, and the neutrons captured predominantly on ³⁵Cl nuclei, which have a much larger neutron capture cross-section than deuterium nuclei, resulting in a higher neutron detection efficiency. Capture on chlorine also released more energy (8.6 MeV) and yielded multiple γ -rays, which aided in identifying neutron events. In the NCD phase, an array of proportional counters (the Neutral Current Detection, or NCD, array) was deployed in the D₂O (Amsbaugh et al. 2007), but some sensitivity remained in the data from the PMT array for neutrons capturing on the D₂O as in the first phase.

Neutrino events observed with the PMT array were selected with the same selection and reconstruction criteria as those in Aharmim et al. (2011a) except the data-set features no high energy cutoff. It consists almost entirely of events below 20 MeV. Table 1 gives the run times and the rate of events consistent with being neutrinos.

3. Burst Events

3.1. Solar Flares

The Homestake solar neutrino experiment reported a small excess of events possibly correlated with large solar flares (Davis 1994, 1996), which provided a motivation for subsequent searches. If these events were associated with pion production in the solar atmosphere then these would produce both electron and muon neutrinos with energies up to approximately 50 MeV. The Kamiokande-II collaboration has reported no excess of events associated with large solar flares (Hirata et al. 1988, 1990). Analysis of the LSD data has also reported no excess of events associated with large solar flares (Aglietta et al. 1991). Neutrinos could also be produced by low energy beta decay due to excitation in the solar atmosphere. Bahcall (1988) has argued that although solar flares could produce neutrinos, they should not be observable in any current detector.

For the coincidence analysis presented here, the solar flare data is taken from HESSI (the High Energy Solar Spectroscopic Imager), which registers radiation ranging from 3 keV to 17 MeV with highly accurate timing. Extensive data-sets are available from (Mission 2009), with data from over 20000 flares – the analysis performed in this paper requires simply the time (t_j), duration (δt_j) and intensity (I_j) of each flare. I_j is defined as the total number of registered counts and varies over 10 orders of magnitude. To speed up the analysis we only include events with intensity $I_j > 10^5$ photon counts, which reduces the number of events to 842. δt_j is typically around 30 minutes, but is also event-dependent. Data from HESSI starts on modified Julian date (MJD) 52330, thus overlapping with the last half of the salt phase and completely with the NCD phase.

Table 1: Run times and event rates for the three phases of SNO. The dates are given in modified Julian date (MJD).

Phase	On (MJD)	Off (MJD)	Low energy cutoff (MeV)	Livetime (days)	Event rate (day^{-1})
D ₂ O	51484	52056	5.0	282.6	9.29
Salt	52116	52879	5.5	385.5	8.15
NCD	53336	54067	6.5	364.4	4.09

3.2. γ -ray Bursters

γ -ray bursters (GRB's) have been known since the 1960's, and an exhaustive review is provided in (Piran 2004). GRB's produce very large fluxes of γ rays over a definite period of time. There are two classes, short and long. Short GRB's emit 90% of the energy within about 100 ms, while long GRB's emit 90% of the energy within about 100 s. Many models for explaining GRB's have been proposed. Some models suggest a comparatively large flux of high energy (GeV to TeV) ν 's, which might be detectable in (e.g.) ICECUBE. Hence it is reasonable to ask if ν 's could be seen by SNO. The answer is almost certainly not, as fluxes of low-energy (MeV) ν 's should be very low, such that no existing or planned detector could see them from cosmological distances (Piran 2004). However, given the uncertainty in the GRB models, it is useful to look for coincidences between GRB's and SNO events. Note that Super-Kamiokande (SK) (Fukuda et al. 2002; Thrane et al. 2009) has already searched for such coincidences, but with no significant effect observed.

The “time of a GRB” is defined as the time at which the burst starts. Since the real nature of GRB's is unknown, there are various possibilities for the timing of the associated ν 's. In this analysis we have chosen two time windows to cover different possible scenarios:

1. Assume that the ν 's are produced before the γ 's (e.g. a hypernova would have a core collapse followed by a visible outburst after some hours.) Based on SN 1987a the time difference could be as much as 3 hours. For this, we search for ν 's in an asymmetrical window $-3 \text{ hrs} < \delta t < 0$.
2. Assume the initial γ 's then trigger a secondary process which emits ν 's. In addition, assuming the ν 's are massive, they would be

slightly delayed due to the travel time. For this, we search for ν 's in an asymmetrical window $0 < \delta t < 3 \text{ hrs}$.

The GRB data for this analysis is taken from Swift (NASA 2006), which is a multi-wavelength observatory dedicated to the study of GRB's, with three instruments that cover γ -ray, X-ray, UV and optical frequencies. The Swift data set starts on MJD 53329, which overlaps completely with the NCD phase, and includes a total of 190 events.

4. Maximum Likelihood Burst Analysis

4.1. Motivation and Assumptions

Both solar flares and GRB's have hugely varying fluxes. Clearly, if a burst event is “strong”, it is more likely to have associated ν 's; some bursts would be (hypothetically) too weak to produce a ν in SNO. We also need to take into account ordinary solar neutrino (“background”) events in SNO. In addition, the duration of solar flares is highly variable, while various different time windows can be used for GRB's. A Maximum Likelihood Burst Analysis (MLBA) can adeptly deal with all the above challenges. For the MLBA, we suppose the neutrino events in SNO fall into two classes:

1. Random events that arrive at a constant background rate r_B .
2. Burst events that are associated with some astrophysical trigger (i.e. GRB's or solar flares). These consist of n_x events at a time t_j with a characteristic spread δt_j . In the case of a flare, t_j would be the start of the flare and δt_j its duration. In the case of a GRB, t_j and δt_j would be dictated by the time window being used (see Section 3.2).

The simplest assumption is that the number of extra neutrino events in SNO (i.e. burst-associated

events rather than background) is directly related to the fluence:

$$n_x(I) = \alpha I. \quad (1)$$

This assumption would be exact if all bursts have identical physical causes but are at varying distances, so that the variations in both ν and γ fluxes are purely geometric. It is also likely to be approximately true in cases where there is a significant variation in intrinsic luminosity, as in the case of Type II supernovae.

A Maximum Likelihood fit can be used to estimate α by averaging over many bursts. It is well known that supernovae fall into two very different classes: Type Ia (carbon detonation) supernovae produce very few prompt neutrinos (Odrzywolek, A. & Plewa, T. 2011), whereas Type II (core collapse) supernovae emit approximately 10^{57} ν 's over a 10 s period. It is quite possible that GRB's follow a similar pattern. In such a case it should be noted that we have fitted for the average α over all bursts, and if only some bursts include neutrino emissions then the α for these bursts would be higher.

4.2. Maximum Likelihood Fitting

We have a total N_{SNO} events spread over a total time T_{tot} and a live time T_{live} . Assuming any actual signal is small, we have a background rate

$$r_B = \frac{N_{\text{SNO}}}{T_{\text{live}}} \quad (2)$$

Note that the background rates r_B for the various phases are given in Table 1. Now suppose there is a burst with some characteristic intensity I_j at time t_j which lasts for a time δt_j . By hypothesis, this will produce $n_x(I_j) = \alpha I_j$ ν 's. In time δt_j around the j 'th burst, we would expect

$$n_j = w_j (r_B \delta t_j + \alpha I_j) \quad (3)$$

events, where w_j is a weighting factor for the detector livetime given by

$$w_j = \frac{1}{\delta t_j} \int_{t_j}^{t_j + \delta t_j} \sum_m H(t - t_m^s) H(t_m^e - t) dt \quad (4)$$

where t_m^s (t_m^e) are the start (end) times of run m , and $H(x)$ is the Heaviside function. If $w_j < 0.05$

the burst is ignored altogether. This implicitly assumes that the events are uniformly distributed over the time window. The actual number of events will form a Poisson distribution, so we would observe k_j events with a relative probability

$$P(k_j, \alpha) = \frac{n_j^{k_j}}{k_j!} e^{-n_j} \quad (5)$$

Hence for all the bursts taken together, the likelihood is:

$$L(\alpha) = \prod_{j=1}^{N_{\text{bursts}}} P(k_j, \alpha) \quad (6)$$

The best-fit value, α_{fit} , is found by minimizing $-\ln(L)$ with respect to α . The 90% upper limits on α are given by

$$-\ln(L(\alpha_{90})) = -\ln(L(\alpha_{\text{fit}})) + 0.821762. \quad (7)$$

It is possible for the fit to give $\alpha_{\text{fit}} < 0$, which is obviously unphysical, but in such a case Eq (7) can still give us a physical upper limit on α .

An upper limit can also be obtained via Monte Carlo simulations using a Feldman-Cousins style (Feldman & Cousins 1998) method, which provides a completely independent check on the method. This involves creating many MC datasets with some fixed α value (α_{in}) and running each dataset through the Maximum Likelihood fitter to obtain a result α_{out} , repeating this procedure for many different α_{in} values, then plotting the α_{in} vs α_{out} distributions. We can extract from this an upper limit on α_{in} (i.e. the ‘‘real’’ value of α) for a given value of α_{out} (i.e. the value of α obtained from the Maximum Likelihood fitter). Results for this technique are completely consistent with the MLBA results reported herein.

4.3. Obtaining Fluence Limits

Once a limit on the number of burst-related SNO events is obtained, the next logical step is to turn this into a fluence limit. Due to the lack of well known spectra, previous results documented in the literature (Fukuda et al. 2002; Hirata et al. 1988) express limits on neutrino fluences at the detector in terms of ‘‘Green’s function’’ fluence for

mono-energetic neutrinos:

$$\Phi_\nu(E_\nu) = \frac{N_{90}}{\sum_{i=1}^{n_R} N_i \int \sigma_i(E'_\nu) \varepsilon_i(E'_\nu) \delta(E_\nu - E'_\nu) dE'_\nu}. \quad (8)$$

This form takes into account the possibility that there may be several (n_R) possible reactions for detecting any particular neutrino flavour. For a given reaction i , $\sigma_i(E_\nu)$ is the cross-section for neutrinos of energy E_ν , $\varepsilon_i(E_\nu)$ the detection efficiency, N_i the total number of targets and N_{90} the 90-percent confidence limit on the number of burst-associated neutrinos registered at the detector. Note that only data from the photomultiplier array in the SNO detector was used, not the data from the independent array of neutron detectors in the NCD phase of the experiment.

To calculate N_{90} in Eq (8), we start by taking the fitted value α_{90} (the 90% confidence limit for α .) From this, we find the expected number of burst-related ν 's per burst from Eq (1), weighted by the windowing function in Eq (4). Altogether, this yields:

$$N_{90} = \frac{\alpha_{90}}{N_{burst}} \sum_{j=1}^{N_{burst}} I_j w_j \quad (9)$$

Note that this does not include systematic uncertainties, which are small compared to the statistical uncertainties.

For each of the reactions we parameterize the low-energy cross-section in Eq (8) as

$$\sigma_i(E) = \sigma_i^0 (E_\nu - E_i^0)^{k_i} \quad (10)$$

where the values of the threshold energy E_i^0 are calculated, and σ_i^0 and k_i are fitted using energies below 20 MeV. Each reaction also has its own $\varepsilon_i(E_\nu)$, which depends on one or more factors (photomultiplier tube efficiency for electron detection, neutron capture efficiency, etc.) and varies from phase to phase in the SNO experiment. The number of targets also varies from reaction to reaction.

5. Results

5.1. Flare Results

The MLBA results for α for the solar flares are shown in Table 2: the larger number of flares during the SNO salt phase is due to the overlap with

the solar maximum. There is no evidence for a signal, for either the SNO salt phase or NCD phase.

Figure 2 shows a comparison of the excess observed in Homestake run 117 in comparison to exclusion limits from the results presented here and KAM II (Hirata et al. 1988). Excesses were claimed to be seen in other Homestake runs with similar limits. The results are obtained assuming all the neutrinos are generated as ν_e 's in the Sun's atmosphere, but due to vacuum oscillations the probability of detecting the neutrinos in that state is 0.55. The remaining probability has the neutrinos as either ν_μ or ν_τ . This analysis excludes the Homestake result down to approximately 2.2 MeV. The KAM II and Homestake results are both obtained from single large bursts, whereas this result is obtained from multiple bursts. Also the Homestake result is obtained with a burst occurring during a run that lasted six days (MJD 48408.89 to 48414.85), whereas this result is obtained with a significantly shorter interval corresponding to the actual flare.




Fig. 2.— Fluence of neutrinos from solar flares versus neutrino energy. The excess shown for Homestake occurred during run 117, which corresponded to a large solar flare. The Homestake result is obtained assuming the excess in run 117 is attributed to a solar flare. The results for all experiments are calculated assuming pure ν_e production in the solar atmosphere and include vacuum oscillations during their journey to Earth. The results for Kam II are recalculated from (Hirata et al. 1988) based on this model and assuming a 100% detection efficiency for scattered electrons above 19 MeV and 0% efficiency below this.

Table 2: MLBA results for solar flares. Note that the units of α are [SNO events/HESSI photon counts]. α_{90} refers to the 90% upper confidence bound on α , and N_{90} is the 90% upper confidence bound on the average number of flare-associated SNO events per flare.

SNO Data	α_{fit}	α_{90}	$\ln[L(\alpha)] - \ln[L(0)]$	No. Flares	N_{90}
Salt Phase	2.79×10^{-9}	1.21×10^{-8}	0.13	172	0.057
NCD Phase	5.38×10^{-10}	3.44×10^{-9}	0.07	94	0.022

5.2. GRB Results

The MLBA results for the GRB’s are shown in Table 3. There is no evidence for a signal in either timing window. The 90% upper confidence bound on α obtained from the Maximum Likelihood fitting was verified using Monte Carlo simulation and the Feldman-Cousins technique as described at the end of Section 4.2.

To facilitate comparison with the results of SK (Fukuda et al. 2002), Table 4 converts the Swift results for the NCD phase ($-3 \text{ hrs} < \delta t < 0$ timing window) to a “Green’s function” fluence, using Monte Carlo to obtain the necessary upper limits on the number of GRB-associated events. These results are plotted in comparison to SK’s in Figure 3 for various neutrino flavours.




Fig. 3.— Fluence of neutrinos from GRB versus neutrino energy. The results for SK are obtained from (Fukuda et al. 2002). The authors do not calculate results below 7 MeV, but the limits are expected to have a sharp turn up due to the threshold for detecting scattered electrons. The results from SNO presented here are from the NCD phase using SWIFT data with a $-3 \text{ hrs} < \delta t < 0 \text{ hrs}$ time window. Results from other time windows and phases are similar.

6. Other Events

Two isolated astrophysical events are worth examining. The first of these events is the “Parkes Burst” (Lorimer et al. 2007), a very intense radio signal with a duration of less than 5 ms found in archival data from the Parkes radio telescope. It occurred on 24 August 2001 at UT 19:50:01 (MJD 52145.82640046). The source is unknown, but is believed to be at great distance. The second is a giant γ -ray flare from the magnetar SGR 1806-20 (Palmer et al. 2005) on 27 December 2004 at UT 21:30:26.65 (MJD 53366.89614178). In this case, there was a precursor burst 143 s before the main burst.

As well as searching in the SNO neutrino dataset employed in the rest of this paper, we have also searched a different set of SNO events: the “muon” set employed in (Aharmim et al. 2009). The “muon” set, comprised of 77285 events, is designed to enhance the muon signal rather than the neutrino signal; approximately 500 of the events are likely to be atmospheric neutrinos while the rest are likely muons. By searching both SNO data-sets, we can look for correlations of either neutrinos or muons with these isolated astrophysical events. (High energy muon-neutrinos from any astrophysical source would produce muons in the rock, which is confirmed by the observation of upward going muons. It is straightforward to test for SNO muon events that are time-correlated with one of these isolated astrophysical events.)

We have searched for anomalous SNO events in short time-windows ($\pm 180 \text{ s}$) around these two astrophysical events: results are shown in Table 5. Obviously there is no effect, and the SNO event rate at these times is compatible with the background rate.

Table 3: MLBA results for GRB’s, for the Swift dataset (with the SNO NCD data). Note that units of α are [SNO events \times cm²/10⁻⁷ erg]. N_{90} is the 90% upper confidence bound on the number of GRB-associated SNO events, based on α_{90} .

Timing	α_{fit}	α_{90}	$\ln[L(\alpha_{fit})] - \ln[L(0)]$	No. GRB’s	N_{90}
-3 hrs $< \delta t < 0$	-2.6×10^{-4}	2.1×10^{-3}	0.013	116	0.084
$0 < \delta t < 3$ hrs	1.2×10^{-3}	3.9×10^{-3}	0.286	116	0.164

Table 4: GRB fluence 90% CL upper limits: “Green’s function” fluences, using MLBA results for Swift data with the SNO NCD phase (-3 hrs $< \delta t < 0$ hrs timing window.)

Energy [MeV]	Φ_{ν_e} [cm^{-2}]	$\Phi_{\bar{\nu}_e}$ [cm^{-2}]	$\Phi_{\nu_x}, \Phi_{\bar{\nu}_x}$ [cm^{-2}]
5	3.89×10^{11}	2.79×10^{11}	3.92×10^{11}
7	3.96×10^{10}	3.83×10^{10}	9.64×10^{10}
9	1.92×10^9	4.50×10^9	3.51×10^{10}
11	6.43×10^8	1.44×10^9	1.99×10^{10}
13	4.01×10^8	8.28×10^8	1.35×10^{10}

Table 5: Isolated astrophysical events. The number of SNO events observed is in a ± 180 s window around the event. The number of SNO events expected is calculated based on the rate of “ordinary” SNO neutrino or muon events, which here constitute the “background”. Note that due the lower energy and therefore stricter requirements on backgrounds for the neutrino data compared to the muon data, the neutrino data did not include the run corresponding to SGR 1806.

Astrophysical Event	SNO Data-set	SNO Events Observed	SNO Events Expected
Parkes Burst	Muon	0	2.62
Parkes Burst	Neutrino	3	1.69
SGR 1806	Muon	1	2.62

7. Conclusions

We can conclude that SNO has found no evidence for low energy neutrinos produced in coincidence with solar flares or GRB's. The MLBA technique allows for a robust search for temporal correlations between events recorded in detectors and astrophysical events of widely varying intensities. In addition, regarding neutrino fluences from solar flares, SNO has provided limits in the low-energy ν regime that are improved compared to previous analyses in the literature.

8. Acknowledgments

This research was supported by the following. Canada: Natural Sciences and Engineering Research Council, Industry Canada, National Research Council, Northern Ontario Heritage Fund, Atomic Energy of Canada, Ltd., Ontario Power Generation, High Performance Computing Virtual Laboratory, Canada Foundation for Innovation, Canada Research Chairs; US: Department of Energy, National Energy Research Scientific Computing Center, Alfred P. Sloan Foundation; UK: Science and Technology Facilities Council; Portugal: Fundação para a Ciência e a Tecnologia. We thank the SNO technical staff for their strong contributions. We also wish to thank Dr. Richard Hemingway for his encouragement in this work and Dr. Vicki Kaspi for suggesting SGR 1806 as a candidate.

REFERENCES

- Aglietta, M., Badino, G., Bologna, G., et al. 1991, *Astrophys.J.*, 382, 344
- Aharmim, B., et al. 2005, *Phys.Rev.*, D72, 052010
- . 2009, *Phys.Rev.*, D80, 012001
- . 2010, *Astrophys.J.*, 710, 540
- . 2011a, *arXiv:1109.0763 [nucl-ex]*
- . 2011b, *Astrophys.J.*, 728, 83
- Amsbaugh, J. F., Anaya, J., J.b, B., et al. 2007, *Nucl.Instrum.Meth.*, A579, 1054
- Bahcall, J. N. 1988, *Phys.Rev.Lett.*, 61, 2650
- Boger, J., et al. 2000, *Nucl.Instrum.Meth.*, A449, 172
- Davis, R. 1994, *Prog.Part.Nucl.Phys.*, 32, 13
- . 1996, *Nucl.Phys.Proc.Suppl.*, 48, 284
- Feldman, G. J., & Cousins, R. D. 1998, *Phys.Rev.*, D57, 3873
- Fukuda, S., et al. 2002, *Astrophys.J.*, 578, 317
- Hirata, K., Kajita, T., Kifune, T., et al. 1988, *Phys.Rev.Lett.*, 61, 2653
- Hirata, K. S., et al. 1990, *Astrophys.J.*, 359, 574
- Lorimer, D. R., Bailes, M., McLaughlin, M. A., Narkevic, D. J., & Crawford, F. 2007, *Science*, 318, 777
- Mission, N. G. R. 2009, *HESSI, Electronic Catalogue*, URL: <http://hesperia.gsfc.nasa.gov/hessi/>
- NASA. 2006, *Swift, Electronic Catalogue*, URL: <http://heasarc.gsfc.nasa.gov/docs/swift/swiftsc.html>
- Odrzywolek, A., & Plewa, T. 2011, *Astron.Astrophys*, 529, A156
- Palmer, D. M., Barthelmy, S., Gehrels, N., et al. 2005, *Nature*, 434, 1107
- Piran, T. 2004, *Rev.Mod.Phys.*, 76, 1143
- Thrane, E., et al. 2009, *Astrophys.J.*, 697, 730